

WBA Patient Case Complexity

| | |
|---|---|
| <p>LOW complexity cases include those that are best described as:</p> | <ul style="list-style-type: none"> • A patient with a single-system presentation, with minimal complications (medical and/or social) and responsive to first line treatment. • A patient with a self-evident diagnosis where management is straightforward. • A stable patient, with a common presentation or a clear diagnosis. <p>Modifiers: No modifiers such a language, mental health status, social, representation or inconsistent clinical findings impacting on assessment or management.</p> <p>Examples of low complexity cases:</p> <ul style="list-style-type: none"> • Isolated limb fracture; • Renal colic • DVT; • Cellulitis; • Pneumonia |
| <p>MEDIUM complexity cases include those that are best described as:</p> | <ul style="list-style-type: none"> • A patient with multi-system presentations, and minimal complications (medical and/or social). • A patient with a single-system presentation and multiple or significant complications <p>or;</p> <ul style="list-style-type: none"> • A patient with a single system presentation and multiple or significant co-morbidities <p>or;</p> <ul style="list-style-type: none"> • A patient with a single-system presentation with at least one modifier or • A stable patient, without a clear diagnosis <p>Modifiers: At least one modifier such as language, mental health status, social representation or inconsistent clinical findings impacting on assessment or management.</p> <p>Examples of medium complexity cases:</p> <ul style="list-style-type: none"> • Fracture with nerve/neurovascular compromise; • Syncope/abdominal pain/chest pain with at least one modifier. • STEMI etc. |
| <p>HIGH complexity cases include those that are best described as:</p> | <ul style="list-style-type: none"> • A patient with multi-system problems and multiple/significant complications (medical and/or social). • A patient with multi-system presentation with multiple or significant co-morbidities <p>or;</p> <ul style="list-style-type: none"> • A patient with multi-trauma <p>or;</p> <ul style="list-style-type: none"> • An unstable/deteriorating patient, with an uncommon presentation or without a clear diagnosis. • A patient presenting with a life/limb/sight threatening condition. <p>Modifiers: At least two modifiers such as language mental health status, social, representation or inconsistent, clinical findings impacting on assessment or management.</p> <p>Examples of high complexity cases:</p> <ul style="list-style-type: none"> • Elderly patient with fracture of secondary to syncope on oral anticoagulants; • Undifferentiated shock; • Immunocompromised patient with shortness of breath with renal failure; • GI bleed patient with chest pain on warfarin with mechanical valve. |

Special patient: e.g. immunosuppressed or compromised, Haemodialysis or renal/other transplant patient, any patient with significant chronic and or complex illness that may impact on current presentation

